



# North Dakota

## Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,\* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
Lowest	45th highest	48th highest

\*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




# Sexually transmitted diseases North Dakota

## Gonorrhea rates, 1982–1992



## Gonorrhea rates, 1992



## National ranking,\* 1992


Males – 48th highest    Females – 48th highest

\*National ranking refers to males and females aged 10–24 years.




## Chlamydia cases and rates,\* 1988–1992


Cases




Rates



Cases



Rates



## Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



\*Chlamydia cases and rates refer to males and females of all ages.

†Missing bar indicates fewer than six cumulative cases.



## Related risk behaviors, 1993 \_\_\_\_\_ North Dakota

*Related risk behavior data are not available for North Dakota.*

*Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data).*

